

The effect of the competitive season in professional basketball on inflammation and iron metabolism

AUTHORS: Dzedzej A¹, Ignatiuk W¹, Jaworska J¹, Grzywacz T², Lipińska P³, Antosiewicz J⁴, Korek A⁵, Ziemann E¹

¹ Gdansk University of Physical Education and Sport, Department of Physiology and Pharmacology, Gdansk, Poland

² Institute of Sport – National Research Institute, Department of Physiology, Warsaw, Poland

³ Institute of Sport – National Research Institute, Department of Biomechanics, Warsaw, Poland

⁴ Medical University of Gdansk, Department of Bioenergetics and Physiology of Exercise, Gdansk, Poland

⁵ Basketball Team, Asseco Prokom, Gdynia, Poland

ABSTRACT: Following acute physical activity, blood hepcidin concentration appears to increase in response to exercise-induced inflammation, but the long-term impact of exercise on hepcidin remains unclear. Here we investigated changes in hepcidin and the inflammation marker interleukin-6 to evaluate professional basketball players' response to a season of training and games. The analysis also included vitamin D (25(OH)D3) assessment, owing to its anti-inflammatory effects. Blood samples were collected for 14 players and 10 control non-athletes prior to and after the 8-month competitive season. Athletes' performance was assessed with the NBA efficiency score. At the baseline hepcidin correlated with blood ferritin ($r=0.61$; 90% CL ± 0.31), but at the end of the season this correlation was absent. Compared with the control subjects, athletes experienced clear large increases in hepcidin (50%; 90% CI 15-96%) and interleukin-6 (77%; 90% CI 35-131%) and a clear small decrease in vitamin D (-12%; 90% CI -20 to -3%) at the season completion. Correlations between change scores of these variables were unclear ($r = -0.21$ to 0.24 , 90% CL ± 0.5), but their uncertainty generally excluded strong relationships. Athletes were hence concluded to have experienced acute inflammation at the beginning but chronic inflammation at the end of the competitive season. At the same time, the moderate correlation between changes in vitamin D and players' performance ($r=0.43$) was suggestive of its beneficial influence. Maintaining the appropriate concentration of vitamin D is thus necessary for basketball players' performance and efficiency. The assessment of hepcidin has proven to be useful in diagnosing inflammation in response to chronic exercise.

CITATION: Dzedzej A, Ignatiuk W, Jaworska J et al. The effect of the competitive season in professional basketball on inflammation and iron metabolism. *Biol Sport*. 2016;33(3):223–229.

Received: 2015-09-07; Reviewed: 2015-11-20; Re-submitted: 2016-01-04; Accepted: 2016-01-14; Published: 2016-05-10.

Corresponding author:

Ewa Ziemann

Gdansk University of Physical Education and Sport

Kazimierza Górskiego 1

80-336 Gdańsk, Poland

Phone: 48585547483

Fax: 48585520751

E-mail: ziemann.ewa@gmail.com

Key words:

Hepcidin

Ferritin

Vitamin D

Interleukin 6

Performance

INTRODUCTION

Effective energy metabolism and transport of oxygen both need to engage several iron-containing proteins, such as cytochromes in the mitochondria, myoglobin and haemoglobin. Thus, proper iron homeostasis is essential, particularly during long lasting athletic competitions. In another way, iron overload can stimulate formation of reactive oxygen species (ROS) [1]. Most of the iron stored is located in ferritin, yet it does not stimulate ROS formation. Still, certain stressors such as UV light, TNF- α or hydrogen peroxide can trigger iron release from ferritin, hence stimulating iron-dependent ROS formation [2]. At the same time, a systemic inflammatory response may be triggered by excessive exercise, common in professional sport [3]. The pro-inflammatory response has the potential to disturb the intracellular iron homeostasis. In consequence, the inflammation acts as a signal activating regeneration processes including proliferation of satellite cells and the molecular mechanism of protein

biosynthesis. Hence, inflammation may contribute to the improvement of adaptation to exercise [4].

Nevertheless, the mechanism responsible for this anti-inflammatory response in training and regular exercise needs to be recognized. Zembron-Lacny and co-workers reported significant shifts in markers of ROS activity – thiol redox status and inflammatory mediators (IL-6, TNF α) – following professional basketball training and found the changes in thiol redox and cytokine response to be correlated [5]. In our previous paper we concluded that the pro-inflammatory cytokine IL-1 β , most likely induced by an extensive workload during the tournament season, was responsible for the low level of ferritin in young, professional tennis players [6]. Moreover, the study revealed that increased levels of the pro-inflammatory cytokines were associated with an elevated level of hepcidin – a liver-produced hormone regulating iron metabolism in the gut and macrophages [7, 8]. These

data are in agreement with the paper by Nemeth, who stated that the inflammatory cytokine interleukin-6 (IL-6) is the primary mediator responsible for regulating hepcidin activity [9]. Hepcidin is also expressed in response to hypoxia, elevated iron levels and exercise [10, 11]. Clinical research demonstrated the peak hepcidin level to occur 3 h following the IL-6 peak rise [12]. The relationship between hepcidin and IL-6 was confirmed, investigating the response to a single dose of exercise: endurance [13] as well as interval training [14]. In addition, Roecker and co-workers observed a considerable increase in hepcidin level 24 h after a 42.2-km marathon run in female athletes [15]. Data determining the elevated resting serum hepcidin values in response to long-lasting training and competition in professional athletes are, however, limited. Ma and co-workers showed that hepcidin was not chronically elevated with sustained training in competitive collegiate runners [16]. Still, 85% of those runners were supplemented with iron. A significant elevation in hepcidin was also observed in specific groups with anaemia, associated with inflammation and/or chronic diseases [17].

These data confirm that inflammation plays a significant role in hepcidin synthesis. In physical training, especially when combined with competition stress, it is essential to maintain the appropriate balance between the pro- and anti-inflammatory response [3]. We assumed that changes in iron status and hepcidin response may be one of the mechanisms which helps in achieving this goal.

Therefore, the first aim of our study was to evaluate hepcidin level among professional basketball players at baseline and in response to long-lasting physical workload, as well as training and a game. Owing to the well-documented anti-inflammatory effect of vitamin D (25(OH)D3) [18], we assumed that lower hepcidin would be observed among athletes with higher levels of vitamin D.

MATERIALS AND METHODS

Data collection. To monitor the influence of training and game workload on iron status, blood analysis was conducted in a professional basketball team. The same blood parameters were assessed in the control group as well. Body composition was also measured. These examinations were performed at the onset of the season (September, baseline) and two days directly after the last game of the first round of the games (March). This 8-month period in-between included all indoor games scheduled by the premier basketball league. Throughout this period physical training and game workload as well as each game efficiency were recorded. All the subjects were informed of the aim of the study and gave their written consent for participation in the project. The protocol of the study was approved by the local ethics committee, in accordance with the Declaration of Helsinki.

Subjects

Two groups took part in the investigation: basketball players and non-athletes (as a control group). On average, the players had 20 ± 5.0 years of training and competition experience. Non-athletes were

instructed not to change their life style in the period between examinations.

Body composition assessment

Body mass (BM) and body composition were estimated using a multi-frequency impedance plethysmograph body composition analyser (InBody 720, Biospace, Korea). Using a diverse range of frequencies from 1 kHz to 1 MHz, InBody 720 accurately measured the amount of body water as well as body composition, including fat mass, free fat mass and skeletal muscle mass. Precision of the repeated measurements was expressed as the coefficient of variation, which on average was 0.6% for the fat mass percentage [19, 20]. The measurements were taken one hour before breakfast. The participants had emptied their bladders and bowels prior to the assessment. During the measurement, the participants wore only briefs and remained barefoot [21].

Training and game workload

The training workload during this period was recorded by coaches. The team analyst recorded the NBA efficiency score for each player in each game as a measure of players' performance. The formula created by John Hollinger and modified by the statistician Martin Manley is officially used by NBA and Basketball-Reference.com. The change in the score for each player over the season was derived by regressing the score against the game number (28 games) and deriving the difference between predicted scores in the first and last games (the slope times 27).

Blood sampling and cytokine analysis

Blood samples were taken from an antecubital vein into single-use containers with an anticoagulant (EDTA_{K2}). After collection, the samples were immediately stored at a temperature of 4°C. Within 10 minutes, they were centrifuged at 3000 g and 4°C for 10 min. Aliquots of the plasma were stored at -80°C.

Blood was collected at rest, at the same time of the day (8.00 a.m.). Red blood cell counts [$10^6 \mu\text{L}$] (RBC), haematocrit [%] (Hct) and blood haemoglobin concentration [$\text{g} \cdot \text{dL}^{-1}$] (Hb) were determined from the venous blood samples by conventional methods using a COULTER LH 750 Hematology Analyser (Beckman-Coulter, USA). Plasma interleukin-6 (IL-6) level was determined via the enzyme immunoassay method using commercial kits (R&D Systems, USA). The detection limit for IL-6 was $0.039 \text{ pg} \cdot \text{mL}^{-1}$, with the average intra-assay CV below 8.0%.

Serum levels of 25(OH)D3 were determined by electrochemiluminescence (ECLIA) using the Elecsys system (Roche, Switzerland). The intra assay and inter assay coefficients of variation (CVs) were 5.6% and 8.0%, respectively, and the limit of detection was $4 \text{ ng} \cdot \text{mL}^{-1}$.

Serum hepcidin was determined using the DRG Elisa kit (DRG Instruments, GmbH, Germany) according to the manufacturer's protocol. This assay detects the 25 amino-acid form of hepcidin – the

biologically active form of the hormone. The dynamic range of the assay was 0.9-140 ng·mL⁻¹. The average CVs precisions of intra-assay was 4.41% and 9.70%, respectively. The normal 5–95% range in healthy adults described by the manufacturer corresponds to 13.3–54.4 ng·mL⁻¹. Analytical sensitivity of the hepcidin kit was calculated by subtracting two standard deviations from the mean of 20 replicate analyses of the zero standard. Similar hepcidin measurements (Elisa kit) were used in a previous study [22] – at the moment it seems to be the most time- and cost-effective method.

Statistical analysis

Statistical analysis was performed in two ways. Comparison of body composition and its changes was performed using STATISTICA 9.0 (StatSoft, USA). Results are expressed as the mean and standard deviation (X±SD). The normality of data was tested using the Shapiro-Wilks W-test. The level of significance was set at 0.05 for all analyses. Additionally, to determine the differential significance at baseline and after an 8-month period the t-test was used. Measures related to haematological parameters were analysed with a spreadsheet for analysis of parallel-group controlled trials, and effects were interpreted using magnitude-based inferences [23,24]. All data were log-transformed to reduce bias arising from the error non-uniformity. Means of the score changes in both groups, standard deviations of the score changes and the obtained effects (differences in the changes of the means and their certainty limits) were back-transformed to percentage units. Mean changes and effects were adjusted to the overall mean baseline value for both groups by including the baseline value as a covariate in the analysis. Magnitudes of the effects were evaluated with the log-transformed data by standardising with the standard deviation of the baseline values of the control group. Threshold values for assessing magnitudes of standardised effects were 0.20, 0.60, 1.2 and 2.0 for small, moderate, large and very large respectively. Uncertainty in each effect was expressed as the 90% confidence limit as well as the probability that the true effect was substantially positive (an increase) or negative (a decrease). These probabilities were used to make a qualitative, probabilistic, non-clinical inference about the true effect: if the probability of the effect being a substantial increase or a substantial decrease was >5% in both cases (equivalent of 90% confidence interval overlapping thresholds for a substantial increase and decrease), the effect was reported as unclear; otherwise, it was considered clear and assigned the relevant magnitude value, with the qualitative probability that the true effect was a substantial increase, substantial decrease, or a trivial difference (whichever outcome had the largest probability). The following scale was used for interpreting the probabilities: 25–75%, possible; 75–95%, likely; 95–99.5%, very likely; >99.5%, most likely. This study involved assessment of substantial changes in five measures. To maintain an overall error rate of <5% for declaring one or more changes to have opposite magnitudes (a substantial decrease instead of an increase, and vice versa), the effects were also evaluated as clear or unclear with a threshold of 5%/5 (1%),

equivalent to consideration of overlap of substantial values with a 98% confidence interval.

The spreadsheet [23] also provided an estimate of a standard deviation representing individual responses to the season of training in the athletes (the variation in the change scores in the athlete group additional to that in the control group). The standard deviation was given by the square root of the difference in the variance of the change scores, and confidence limits were provided by assuming asymptotic normality of the variance. When the difference in the variance or a confidence limit was negative, the sign was changed before taking the square root, and the standard deviation was shown as a negative value. Individual responses were interpreted with magnitude-based inference as above, but the standardised magnitude thresholds were halved to effectively assess the magnitude of twice the standard deviation. The usual scale for correlation coefficients (0.1, 0.3, 0.5, 0.7 and 0.9 for low, moderate, high, very high and nearly perfect, respectively) was used for interpretation of relationships between selected variables and players' performance. Confidence limits (90% CL) for coefficients of correlation were calculated using dedicated spreadsheet.

RESULTS

All basketball players completed the tournament round with no adverse events or injury (serious, excluded from the games) being reported. Athletes' body compositions are outlined in Table 1, which shows only significant changes in water status (intracellular and extracellular water amount) that took place during the investigated period. Changes in body composition for non-athletes were not significant. In agreement with our expectations, non-athletes differed in comparison to basketball players (data for non-athletes are not presented).

Baseline values of basic morphological measures from the athlete and control groups are presented in Table 2. In order to evaluate the influence of the tournament season on changes in blood parameters the same analyses were repeated. Obtained data are also presented in Table 2, including changes in basic measures in both groups throughout the season and a comparison of the observed changes.

TABLE 1. Anthropometric characteristics of basketball players.

Variable	Baseline	After season	P value
Height (cm)	196.7±8.7	196.7 ±8.7	ns
Weight (kg)	94.4 ±8.0	92.1 ±6.4	ns
FFM (kg)	84.7 ±8.7	83.4 ±7.0	ns
SMM (kg)	49.1 ±5.2	48.2 ±4.2	ns
Fat (kg)	9.3 ±3.7	8.7 ±2.5	ns
Fat (%)	9.8 ±4.0	9.5 ±3.0	ns
VFA (cm ²)	57.8 ±25.4	62.9 ±15.5	ns
ECW (kg)	23.2 ±2.5	22.9 ±2.5	0.002
ICFW (kg)	39.1 ±4.2	39.0 ±4.1	ns

Note: Values are means ± SD; FFM – fat-free mass, SMM – skeletal muscle mass, Fat – fat mass, Fat% – percentage of body fat, VFA – visceral fat area, ns – no statistically significant differences between measurements.

TABLE 2. Basic measures at baseline and changes in the measures across a season in basketball players and non-athlete controls.

		Baseline	Observed change	Observed effect	Inference
		mean \pm SD	mean \pm SD	mean; CI	
WBC ($\times 10^3 \cdot \mu\text{L}^{-1}$)	Athletes	6.5 \pm 0.7	7 \pm 23 %	2%;	trivial, unclear
	Controls	5.9 \pm 0.9	6 \pm 20 %	-11 to 17%	
RBC ($\times 10^3 \cdot \mu\text{L}^{-1}$)	Athletes	4.92 \pm 0.29	1.5 \pm 3.9 %	2.0%;	trivial, unclear
	Controls	4.94 \pm 0.34	-0.4 \pm 7.0 %	-2.2 to 6.4 %	
HGB (g·dL ⁻¹)	Athletes	14.9 \pm 0.6	3.5 \pm 3.9 %	6.9%;	moderate \uparrow ****
	Controls	15.2 \pm 0.9	-3.1 \pm 4.6 %	3.7 to 10.2 %	
Hct (%)	Athletes	43.0 \pm 1.6	2.2 \pm 3.7 %	4.6%;	moderate \uparrow ****
	Controls	44.2 \pm 2.0	-2.3 \pm 2.3 %	2.5 to 6.8 %	
MCV (fL)	Athletes	87.5 \pm 3.6	0.8 \pm 1.5 %	2.9%	small \downarrow *
	Controls	90.0 \pm 5.4	-0.2 \pm 8.0 %	-1.7 to 7.6 %	
MCH (pg)	Athletes	30.3 \pm 1.3	2.1 \pm 2.1 %	5.2%	moderate \uparrow ***
	Controls	30.9 \pm 2.1	-2.9 \pm 9.2 %	-0.2 to 10.8 %	
MCHC (g·dL ⁻¹)	Athletes	34.6 \pm 0.7	1.3 \pm 2.4 %	2.2%	moderate \uparrow **
	Controls	34.3 \pm 0.8	-0.9 \pm 3.7 %	-0.2 to 4.6 %	
RDW (%)	Athletes	13.0 \pm 0.4	0.4 \pm 2.6 %	0.3%	trivial, unclear
	Controls	13.1 \pm 0.5	0.1 \pm 5.0 %	-2.7 to 3.4 %	
PLT ($\times 10^3 \cdot \mu\text{L}^{-1}$)	Athletes	274 \pm 29	-4 \pm 19 %	1%	trivial, unclear
	Controls	238 \pm 43	-4 \pm 30 %	-15 to 19 %	
MPV (fL)	Athletes	8.6 \pm 0.8	1.3 \pm 6.3%	-2.0%	small \downarrow , unclear
	Controls	10.5 \pm 0.7	3.3 \pm 10.4 %	-8.0 to 4.3 %	

Note: CI – 90% confidence interval.; \uparrow increase; \downarrow , decrease. All data are percentages, with the exception of baseline values expressed in measurement units. Likelihood that the true effect is substantial: * – possible, ** – likely, *** – very likely, **** – most likely.

TABLE 3. Measures related to iron metabolism at baseline and changes in the measures across a season in basketball players and non-athlete controls.

		Baseline	Observed change	Adjusted change ^a	Adjusted effect ^b	Inference
		mean \pm SD	mean \pm SD	mean \pm SD	mean; CI	
Hepcidin (ng·mL ⁻¹)	Athletes	69 \pm 14	25 \pm 32 %	71 \pm 25 %	50%;	large \uparrow ***
	Controls	30 \pm 7	44 \pm 18 %	14 \pm 14 %	15 to 96 %	
IL-6 (pg·mL ⁻¹)	Athletes	1.14 \pm 0.54	16 \pm 51 %	21 \pm 46 %	77%;	large \uparrow ****
	Controls	0.95 \pm 0.37	-21 \pm 87 %	-32 \pm 42 %	35 to 131 %	
Ferritin (ng·mL ⁻¹)	Athletes	115 \pm 76	-17 \pm 45 %	-15 \pm 40 %	16%;	small \uparrow *
	Controls	93 \pm 54	-24 \pm 29 %	-27 \pm 21 %	-4 to 40 %	
Iron ($\mu\text{g} \cdot \text{dL}^{-1}$)	Athletes	129 \pm 44	-15 \pm 49 %	-18 \pm 33 %	-14%;	small \downarrow **
	Controls	140 \pm 34	-6 \pm 10 %	-5 \pm 10 %	-26 to -1 %	
Vitamin D (ng·mL ⁻¹)	Athletes	25 \pm 7	-27 \pm 12 %	-27 \pm 13 %	-12%	small \downarrow **
					-20 to -3 %	

Note: CI – 90% confidence interval. All data are percentages, with the exception of baseline values expressed in measurement units. Inferences shown in bold are clear at the 98% level of confidence. ^a – Adjusted to overall mean of athletes and controls at baseline. ^b – Adjusted mean change in athletes minus adjusted mean change in controls.

\uparrow – increase; \downarrow – decrease. Likelihood that the true effect is substantial: * – possible, ** – likely, *** – very likely, **** – most likely.

The main purpose of this study was to evaluate whether the whole tournament season impaired iron metabolism and if these changes were related to hepcidin and IL-6 concentrations. This hypothesis was based on previous observations from professional athletes, both basketball and tennis players, who exhibited low ferritin levels ($<25 \mu\text{g} \cdot \text{L}^{-1}$). Ferritin levels in blood should reflect the intracellular ferritin concentration. Cellular ferritin synthesis also changes in

parallel to cellular iron levels. Hence, low blood ferritin is considered to be the best indicator of iron deficiency. Nonetheless, contrarily to our assumptions the average values of parameters characterising the iron status in the athlete group were in their reference ranges at the baseline as well as at the game round completion. Table 3 shows baseline values of the measures related to iron status in the athlete and control groups.




FIG. 1. The relationship between change scores and baseline values in the athlete and control groups for one of the measures, hepcidin. The figure shows how the mean changes and their differences were adjusted to the mean baseline hepcidin concentration: the adjusted changes in each group are the changes predicted for the mean baseline value using the linear regression.

The baseline concentration of hepcidin in the athlete group was over twice that of the control subjects (a difference of 129%; 90% CI 95 to 168%; very large), while that for vitamin D was lower (-21%; 90% CI -35 to -5%; moderate). Differences for IL-6 (18%; 90% CI -9 to 53%), ferritin (18%; 90% CI -22 to 79%) and iron (-11%; 90% CI -28 to 11%) were small but unclear. The changes in the measures across an 8-month period in each group and the comparison of the changes adjusted to the overall mean at baseline are also shown in Table 3. There were large, clear increases in hepcidin and IL-6, and a small increase in ferritin. Iron and vitamin D exhibited small clear reductions. The effects for hepcidin, IL-6 and vitamin D were still clear at the 98% level. Moreover, a strong correlation between ferritin and hepcidin concentration at the baseline was recorded ($r=0.61$; 90%CL ± 0.31) at the baseline among athletes. This relationship was maintained at the end of the season, but at that time it became only moderate ($r=0.43$; 90%CL ± 0.39). In addition, the relationship between change scores and baseline values in the athletes and control groups for one of the measures, hepcidin, is illustrated in Figure 1. It shows how the mean changes and their differences were adjusted to the mean baseline hepcidin concentration: the adjusted changes in each group are the changes predicted for the mean baseline value using the linear regression.

Individual responses expressed as coefficients of variation were clear for iron (31%; 90% CI 14 to 43%; large) and for ferritin (32%; 90% CI -6 to 49%; moderate). Individual responses for hepcidin (20%; 90% CI -5 to 30%) and IL-6 (16%; 90% CI -29 to 49%) had respectively large and moderate observed values but were unclear, while those for vitamin D (2%; 90% CI -12 to 13 %) were trivial and unclear. The average players' NBA efficiency of all the 28 games was 7.5 with between-athletes SD = 4.5, and within-athletes SD = 5.6. The value of average slope of efficiency was -0.1 ± 3.3 . A likely

positive tendency was noted between change in vitamin D and change in players' performance measure over the season ($r=0.43$; 90%CL ± 0.41). Correlations between changes in hepcidin and IL6 vs changes in vitamin D in the athlete group were $r=0.40$; 90%CL ± 0.42 and 0.20 ; 90%CL ± 0.46 , whereas in the control group $r=0.23$; 90%CL ± 0.44 and $r= -0,19$; 90%CL ± 0.46 , respectively.

DISCUSSION

In the present study we observed that the tournament season in basketball induced significant changes in iron metabolism in professional players. The recorded rise in blood hepcidin was the most pronounced change. Comparison between changes in blood iron in both groups supports this observation as the adjusted effect showed a small clear decrease.

As mentioned above, the main function of hepcidin is to reduce blood iron availability by inhibiting intestinal iron absorption [10]. Thus, the rise in blood hepcidin usually leads to a drop in blood iron. Moreover, by blocking ferroportin protein, which is an iron exporter, hepcidin can also significantly reduce iron trafficking between different cells such as macrophages, liver cells and many others where the protein ferroportin is present [25]. Iron is the main factor which stimulates hepcidin biosynthesis – an increase in blood iron causes hepatocytes and possibly other cells such as adipocytes and pancreatic cells to liberate hepcidin [26]. In the present study we observed a significant positive correlation between blood ferritin and hepcidin concentration at the baseline. Blood ferritin level is considered to be the best indicator of body iron stores [27]. Thus, the correlation between these two parameters seems to reflect the physiological situation where iron induced changes to limit its accumulation. Interestingly, such a correlation was not observed in athletes after the tournament season, indicating that other factors than iron played a role in hepcidin biosynthesis regulation.

We hypothesized that inflammation could be this factor. It was previously shown that basketball players experience low grade systemic inflammation at the end of the game season [5]. This observation was confirmed in this study. We observed that resting IL-6 rose after the season and IL-6 positively correlated with hepcidin despite the fact that such a correlation had not been noted at the beginning of the season. Moreover, in the control group IL-6 was observed to have dropped, and the comparison between changes in both groups showed a large clear increase, which confirms the adverse effects of the tournament season.

Overall, we propose that uncoupling blood hepcidin and ferritin levels could be a good indicator of the inflammatory process in professional athletes. The biological significance of an increase in blood hepcidin due to inflammation is related to the pro-inflammatory effects of iron [28]. Consequently, iron is described as a molecule capable of activating transcriptional nuclear factor- κ B (NF- κ B) [29]. It might lead to both enhanced gene expression and protein synthesis [30]. Among others, pro-inflammatory mediators such as IL-1 β , IL-6 and TNF- α are all up-regulated by NF- κ B. Reduced iron absorp-

tion due to rising blood hepcidin is one way which allows the inflammatory process to self-limit. It is important to note that the observed changes in iron metabolism seem to have been a very sensitive marker of inflammation in athletes, who probably experienced overreaching from the game season.

One reason why training and game workload induced inflammation could be a low concentration of vitamin D. The anti-inflammatory effects of vitamin D are well-documented [31, 32] the deficiency is often observed in indoor athletes [33]. Moreover, a paper by Kopec et al. showed that even in outdoor sport the recommended levels of 25(OH)D3 were recorded in 50% of the players after the summer period but only in 16.7% of the players after the winter time [34]. Additionally, hypovitaminosis D was also recorded among Tunisian athletes despite living in a sun-rich environment [35]. In our study, the first assessment was performed directly after summer (a period of high exposure to UVB), but the values of vitamin D were low in both groups. The vitamin D deficiency was found to have progressed throughout the season, which could be one of the causes of inflammation. Data confirmed that athletes characterized by a smaller reduction in vitamin D were able to maintain their best performance efficiency during the whole game season. The literature reports that 25(OH)D3 concentration can positively affect muscle function [36] as well as muscle tissue [37]. Agergaard et al. revealed that 12 weeks of resistance training supported by supplementation of vitamin D does not influence the whole muscle hypertrophy or muscle strength, but improved muscle quality in the elderly and fibre type morphology in the young, indicating an effect of vitamin D on skeletal muscle remodelling [38]. Although we did not measure muscle strength, we can speculate based on the obtained correlation that the players characterized by elevated vitamin D exhibited higher exercise tolerance and underwent the recovery process faster, which helped them maintain better performance during the whole investigation period.

In our observations, we noted a parallel increase of the exercise-induced inflammatory markers hepcidin and IL-6. Both markers could be considered as indicators of the inflammatory response in professional athletes. It appears particularly interesting to uncouple the relationship between ferritin and hepcidin at the end of the season, which could also be considered a good marker of inflammation.

CONCLUSIONS

In summary, our current data show that overreaching, which is an effect of the basketball tournament season, leads to an increase in blood hepcidin. The strong correlation observed between blood ferritin and hepcidin faded at the end of the season, indicating that it could be a good and sensitive marker of inflammation and overreaching among athletes. Regular assessment of vitamin D should also be included due to its likely role in ameliorating exercise-induced inflammation and the possible benefit to athletes' performance.

Acknowledgements

The experiment was funded by knowledge grant of Polish Ministry of Science and Higher Education No 0016/RS2/2013/52. We would like to express our thanks to whole team and all students for their engagement in the experiment. We thank Marek Adamski and Cole Hairston, team coaches, for help in conducting the research. Moreover we would like to thank for the medical staff, especially the nurse Anna Szydłowska for her help in blood collection. There are no professional relationships between any of the authors and manufacturers of equipment utilized in this study.

Conflict of interests: the authors declared no conflict of interests regarding the publication of this manuscript.

REFERENCES

1. Yuan X, Cong Y, Hao J, Shan Y, Zhao Z, Wang S, Chen J. Regulation of lip level and ros formation through interaction of h-ferritin with g-csf receptor. *J Mol Biol.* 2004;339(1):131-144.
2. Antosiewicz J, Ziolkowski W, Kaczor JJ, Herman-Antosiewicz A. Tumor necrosis factor-alpha-induced reactive oxygen species formation is mediated by jnk1-dependent ferritin degradation and elevation of labile iron pool. *Free Radic Biol Med.* 2007;43(2):265-270.
3. Cooper DM, Radom-Aizik S, Schwindt C, Zaldivar F Jr. Dangerous exercise: Lessons learned from dysregulated inflammatory responses to physical activity. *J Appl Physiol* (1985). 2007;103(2):700-709.
4. Paulsen G, Mikkelsen UR, Raastad T, Peake JM. Leucocytes, cytokines and satellite cells: What role do they play in muscle damage and regeneration following eccentric exercise? *Exerc Immunol Rev.* 2012;1842-97.
5. Zembron-Lacny A, Slowinska-Lisowska M, Ziemba A. Integration of the thiol redox status with cytokine response to physical training in professional basketball players. *Physiol Res.* 2010;59(2):239-245.
6. Ziemann E, Kasproicz K, Kasperska A, Zembron-Lacny A, Antosiewicz J, Laskowski R. Do high blood hepcidin concentrations contribute to low ferritin levels in young tennis players at the end of tournament season? *J Sports Sci Med.* 2013;12(2):249-258.
7. Nicolas G, Viatte L, Bennoun M, Beaumont C, Kahn A, Vaulont S. Heparin, a new iron regulatory peptide. *Blood Cells Mol Dis.* 2002;29(3):327-335.
8. Park CH, Valore EV, Waring AJ, Ganz T. Heparin, a urinary antimicrobial peptide synthesized in the liver. *J Biol Chem.* 2001;276(11):7806-7810.
9. Nemeth E, Rivera S, Gabayan V, Keller C, Taudorf S, Pedersen BK, Ganz T. IL-6 mediates hypoferrremia of inflammation by inducing the synthesis of the iron regulatory hormone hepcidin. *J Clin Invest.* 2004;113(9):1271-1276.
10. Nemeth E, Tuttle MS, Powelson J, Vaughn MB, Donovan A, Ward DM, Ganz T, Kaplan J. Heparin regulates cellular iron efflux by binding to ferroportin and inducing its internalization. *Science.* 2004;306(5704):2090-2093.
11. Peeling P. Exercise as a mediator of hepcidin activity in athletes. *Eur J Appl Physiol.* 2010;110(5):877-883.
12. Sim M, Dawson B, Landers G, Swinkels DW, Tjalsma H, Trinder D, Peeling P.

- Effect of exercise modality and intensity on post-exercise interleukin-6 and hepcidin levels. *Int J Sport Nutr Exerc Metab.* 2013;23(2):178-186.
13. Skarpanska-Stejnborn A, Basta P, Trzeciak J, Szczesniak-Pilaczynska L. Effect of intense physical exercise on hepcidin levels and selected parameters of iron metabolism in rowing athletes. *Eur J Appl Physiol.* 2015;115(2):345-351.
 14. Antosiewicz J, Kaczor JJ, Kasproicz K, Laskowski R, Kujach S, Luszczek M, Radziminski L, Ziemann E. Repeated „all out” interval exercise causes an increase in serum hepcidin concentration in both trained and untrained men. *Cell Immunol.* 2013;283(1-2):12-17.
 15. Roecker L, Meier-Buttermilch R, Brechtel L, Nemeth E, Ganz T. Iron-regulatory protein hepcidin is increased in female athletes after a marathon. *Eur J Appl Physiol.* 2005;95(5-6):569-571.
 16. Ma X, Patterson KJ, Gieschen KM, Bodary PF. Are serum hepcidin levels chronically elevated in collegiate female distance runners? *Int J Sport Nutr Exerc Metab.* 2013;23(5):513-521.
 17. Deschemin JC, Vaulont S. Role of hepcidin in the setting of hypoferrremia during acute inflammation. *PLoS One.* 2013;8(4):e61050.
 18. Mangin M, Sinha R, Fincher K. Inflammation and vitamin d: The infection connection. *Inflamm Res.* 2014;63(10):803-819.
 19. Lim JS, Hwang JS, Lee JA, Kim DH, Park KD, Jeong JS, Cheon GJ. Cross-calibration of multi-frequency bioelectrical impedance analysis with eight-point tactile electrodes and dual-energy x-ray absorptiometry for assessment of body composition in healthy children aged 6-18 years. *Pediatr Int.* 2009;51(2):263-268.
 20. Volgyi E, Tylavsky FA, Lyytikäinen A, Suominen H, Alen M, Cheng S. Assessing body composition with dxa and bioimpedance: Effects of obesity, physical activity, and age. *Obesity (Silver Spring).* 2008;16(3):700-705.
 21. Ziemann E, Olek RA, Grzywacz T, Antosiewicz J, Kujach S, Luszczek M, Smaruj M, Sledziewska E, Laskowski R. Whole-body cryostimulation as an effective method of reducing low-grade inflammation in obese men. *J Physiol Sci.* 2013;63(5):333-343.
 22. Ganz T, Olbina G, Girelli D, Nemeth E, Westerman M. Immunoassay for human serum hepcidin. *Blood.* 2008;112(10):4292-4297.
 23. Hopkins WG. Spreadsheets for analysis of controlled trials with adjustment for a predictor. *Sports Science.* 2006;1046-50.
 24. Hopkins W, Marshall S, Batterham A, Hanin J. Progressive statistics for studies in sports medicine and exercise science. *Med Sci Sports Exerc.* 2009;41(1):3-12.
 25. Delaby C, Pilard N, Goncalves AS, Beaumont C, Canonne-Hergaux F. Presence of the iron exporter ferroportin at the plasma membrane of macrophages is enhanced by iron loading and down-regulated by hepcidin. *Blood.* 2005;106(12):3979-3984.
 26. Rivera S, Nemeth E, Gabayan V, Lopez MA, Farshidi D, Ganz T. Synthetic hepcidin causes rapid dose-dependent hypoferrremia and is concentrated in ferroportin-containing organs. *Blood.* 2005;106(6):2196-2199.
 27. McKinnon EJ, Rossi E, Beilby JP, Trinder D, Olynyk JK. Factors that affect serum levels of ferritin in Australian adults and implications for follow-up. *Clin Gastroenterol Hepatol.* 2014;12(1):101-108 e104.
 28. Kell DB. Iron behaving badly: Inappropriate iron chelation as a major contributor to the aetiology of vascular and other progressive inflammatory and degenerative diseases. *BMC Med Genomics.* 2009;22.
 29. Lin M, Rippe RA, Niemela O, Brittenham G, Tsukamoto H. Role of iron in nf-kappa b activation and cytokine gene expression by rat hepatic macrophages. *Am J Physiol.* 1997;272(6 Pt 1):G1355-1364.
 30. Pham CG, Bubici C, Zazzeroni F, Papa S, Jones J, Alvarez K, Jayawardena S, De Smaele E, Cong R, Beaumont C, Torti FM, Torti SV, Franzoso G. Ferritin heavy chain upregulation by nf-kappab inhibits tnfa-induced apoptosis by suppressing reactive oxygen species. *Cell.* 2004;119(4):529-542.
 31. Bednarek-Skublewska A, Smolen A, Jaroszynski A, Zalaska W, Ksiazek A. Effects of vitamin d3 on selected biochemical parameters of nutritional status, inflammation, and cardiovascular disease in patients undergoing long-term hemodialysis. *Pol Arch Med Wewn.* 2010;120(5):167-174.
 32. Choi M, Park H, Cho S, Lee M. Vitamin D3 supplementation modulates inflammatory responses from the muscle damage induced by high-intensity exercise in sd rats. *Cytokine.* 2013;63(1):27-35.
 33. Bescos Garcia R, Rodriguez Guisado FA. Low levels of vitamin D in professional basketball players after wintertime: Relationship with dietary intake of vitamin D and calcium. *Nutr Hosp.* 2011;26(5):945-951.
 34. Kopec A, Solarz K, Majda F, Slowinska-Lisowska M, Medras M. An evaluation of the levels of vitamin d and bone turnover markers after the summer and winter periods in polish professional soccer players. *J Hum Kinet.* 2013;38135-140.
 35. Sghaier-Ayadi A, Feki M, Ayed IB, Abene O, Fredj MB, Kaabachi K, Chaouachi A. Vitamin D status and determinants of deficiency in non-supplemented athletes during the winter months in Tunisia. *Biol Sport.* 2015;32(4):281-287.
 36. Koundourakis NE, Androulakis NE, Malliaraki N, Margioris AN. Vitamin d and exercise performance in professional soccer players. *PLoS One.* 2014;9(7):e101659.
 37. Hamilton B. Vitamin D and human skeletal muscle. *Scand J Med Sci Sports.* 2010;20(2):182-190.
 38. Agergaard J, Trostrup J, Uth J, Iversen JV, Boesen A, Andersen JL, Schjerling P, Langberg H. Does vitamin-D intake during resistance training improve the skeletal muscle hypertrophic and strength response in young and elderly men? - a randomized controlled trial. *Nutr Metab (Lond).* 2015;1232.